

Dynamic equations for three different qudits in a magnetic field

E. A. Ivanchenko*

*National Science Center “Kharkov Physicotechnical
Institute”, Institute for Theoretical Physics,
Akademicheskaya Str. 1, Kharkov 61108, Ukraine*

(Dated: April 22, 2009)

Abstract

A closed system of equations for the local Bloch vectors and spin correlation functions of three magnetic qudits, which are in an arbitrary, time-dependent, external magnetic field, is obtained using decomplexification of the Liouville-von Neumann equation. The algorithm of the derivation of the dynamic equations is presented. In the basis convenient for the important physical applications structure constants of algebra $\text{su}(2S+1)$ are calculated.

PACS numbers: 03.67.-a, 03.67.Ac, 03.67.Mn

*yevgeny@kipt.kharkov.ua

I. INTRODUCTION

At present it is not known which experiment will lead to the reliable, prototypical quantum computing device. Quantum systems with two states, called qubits, are taken to be the basic unit for quantum information processing. The quantum systems realised on coupled qubits are widely investigated for the purpose of creation of a quantum computer. In the same time there is a possibility to solve this problem, using the systems consisting from coupled qudits (multilevel systems). Qudits possess a number of properties which differ from properties qubits and may have advantages for quantum information processing. For example, two qudits can be entangled more strongly than two qubits [1]. The main purpose of this study is the derivation of dynamic equations for three qudit system, which is placed in a magnetic field, for entanglement study [2] and other applications. The article is organized as follows. Section II contains the model Hamiltonian including the three-particle interactions. In section III the Liouville-von Neumann equation for the matrix density of three qudits in the variable magnetic field is obtained in the Bloch's representation in terms of local vectors and spin correlation functions taking into account three-particle interactions. We describe conservation laws which control numerical calculations effectively. The Appendix describes the Hermitian basis and analytic formulas for structure constants.

II. MODEL HAMILTONIAN

The Hamiltonian of three coupled different qudits (particles with spin $1/2, 1, 3/2, 2, \dots$) in the external ac magnetic field $\mathbf{h} = (h_1, h_2, h_3)$, where h_1, h_2, h_3 are the Cartesian components of the external magnetic field, we will present in a form of decomposition on a complete set orthogonal Hermitian matrices C_α , operating in the Hilbert space of three qudits $H^{S_1} \otimes H^{S_2} \otimes H^{S_3}$ with $(2S_1 + 1)(2S_2 + 1)(2S_3 + 1)$ dimension

$$\hat{H} = \frac{1}{2} h_{\varrho\sigma\tau} C_\varrho^{S_1} \otimes C_\sigma^{S_2} \otimes C_\tau^{S_3}, \quad (1)$$

where S_1, S_2, S_3 are qudit spins, $\otimes$ - denotes a direct product. $C_\varrho^{S_1}, C_\sigma^{S_2}, C_\tau^{S_3}$ are the Hermitian matrices operating in spaces $H^{S_1}, H^{S_2}, H^{S_3}$ accordingly (see Appendix). The functions $h_{\varrho\sigma\tau}$ contain one-, two- and three-qudit interactions [3], [4], [5].

III. DECOMPLEXIFICATION OF THE LIOUVILLE-VON NEUMANN EQUATION

The Liouville-von Neumann equation for the density matrix ρ , describing the dynamics of a 3-qudit system, has the form

$$i\partial_t \rho = [\hat{H}, \rho], \quad \rho(0) = \rho_0, \quad \rho^+ = \rho, \quad \text{Tr } \rho = 1. \quad (2)$$

It is convenient to rewrite the equation Eq. (2) having presented the density matrix ρ as well as the Hamiltonian $\hat{H}$, in the form of decomposition on a complete set orthogonal Hermitian matrices C_α

$$\rho = \frac{1}{c_1(2S_1+1)c_2(2S_2+1)c_3(2S_3+1)} R_{\alpha\beta\gamma} C_\alpha^{S_1} \otimes C_\beta^{S_2} \otimes C_\gamma^{S_3}, \quad (3)$$

where $c_{1,2,3} = \sqrt{\frac{S_{1,2,3}(S_{1,2,3}+1)}{3}}$. We define the three coherence Bloch vectors $R_{m00}, R_{0n0}, R_{00p}$, which are widely used in the theory of magnetic resonance characterize the local properties of the individual qudits,

$$c_1 c_2 c_3 R_{m00} = \text{Tr } \rho C_m^{S_1} \otimes C_0^{S_2} \otimes C_0^{S_3}, \quad (4a)$$

$$c_1 c_2 c_3 R_{0n0} = \text{Tr } \rho C_0^{S_1} \otimes C_n^{S_2} \otimes C_0^{S_3}, \quad (4b)$$

$$c_1 c_2 c_3 R_{00p} = \text{Tr } \rho C_0^{S_1} \otimes C_0^{S_2} \otimes C_p^{S_3}. \quad (4c)$$

while the tensors $R_{mn0}, R_{m0p}, R_{0np}, R_{mnp}$

$$c_1 c_2 c_3 R_{mn0} = \text{Tr } \rho C_m^{S_1} \otimes C_n^{S_2} \otimes C_0^{S_3}, \quad (5a)$$

$$c_1 c_2 c_3 R_{m0p} = \text{Tr } \rho C_m^{S_1} \otimes C_0^{S_2} \otimes C_p^{S_3}, \quad (5b)$$

$$c_1 c_2 c_3 R_{0np} = \text{Tr } \rho C_0^{S_1} \otimes C_n^{S_2} \otimes C_p^{S_3}, \quad (5c)$$

$$c_1 c_2 c_3 R_{mnp} = \text{Tr } \rho C_m^{S_1} \otimes C_n^{S_2} \otimes C_p^{S_3}, \quad (5d)$$

describe the spin correlations. In the formulas Eq. (4), Eq. (5) $C_0^{S_i} \equiv C_{0,z}^{S_i} = \sqrt{\frac{S_i(S_i+1)}{3}} E_{S_i}$, E_{S_i} is the unit matrix in dimension $(2S_i+1)$ and each of the Latin indices designates a set of matrices $C_{k^{S_i}, q^{S_i}; x}^{S_i}, C_{k^{S_i}, q^{S_i}; y}^{S_i}, C_{k^{S_i}; z}^{S_i}$, where $1 \leq k^{S_i} \leq 2S_i$, and $1 \leq q^{S_i} \leq k^{S_i}$ in $C_{k^{S_i}, q^{S_i}; x}^{S_i}, C_{k^{S_i}, q^{S_i}; y}^{S_i}$ and $1 \leq k^{S_i} \leq 2S_i$, in $C_{k,z}^{S_i}$ in steps of 1, $i = 1, 2, 3$.

Let's formulate the linear algorithm of a decomplexification of the Liouville-von Neumann

equation.

1. Insert the Hamiltonian Eq. (1) and the density matrix Eq. (3), decomposed on Hermitian basis into the Liouville-von Neumann equation Eq. (2).
2. Multiply the Liouville-von Neumann equation by all elements of the basis $C_\eta^{S_1} \otimes C_\theta^{S_2} \otimes C_\vartheta^{S_3}$ in turn.
3. Execute operation of a trace taking for each equation.
4. Apply the formula $\text{Tr}(C_\alpha^{S_1} \otimes C_\beta^{S_2} \otimes C_\gamma^{S_3})(C_\epsilon^{S_1} \otimes C_\epsilon^{S_2} \otimes C_\zeta^{S_3})(C_\eta^{S_1} \otimes C_\theta^{S_2} \otimes C_\vartheta^{S_3}) = \text{Tr} C_\alpha^{S_1} C_\epsilon^{S_1} C_\eta^{S_1} \text{Tr} C_\beta^{S_2} C_\epsilon^{S_2} C_\theta^{S_2} \text{Tr} C_\gamma^{S_3} C_\zeta^{S_3} C_\vartheta^{S_3}$.
5. Express a trace from the three matrices through structure constants according to formulas Eqs. (A4,A5).
6. As a result the use of the structure constants symmetry the real terms in each equation are cancelled out, and the purely imaginary terms are duplicated.
7. The imaginary unit i is cancelled.

This algorithm is easy to apply for the system of more than 3 qudits.

In terms of the functions $R_{\alpha\beta\gamma}$ the Liouville-von Neumann equation becomes real and comprises a closed system of first-order differential equations for the local Bloch vectors and spin correlation functions

$$\partial_t R_{m00} = c_2 c_3 e_{jim}^{S_1} (h_{j00} R_{i00} + h_{jk0} R_{ik0} + h_{j0k} R_{i0k} + h_{jkr} R_{ikr}), \quad (6a)$$

$$\partial_t R_{0n0} = c_1 c_3 e_{ikn}^{S_2} (h_{0i0} R_{0k0} + h_{ji0} R_{jk0} + h_{0ij} R_{0kj} + h_{jir} R_{jkr}), \quad (6b)$$

$$\partial_t R_{00p} = c_1 c_2 e_{ijp}^{S_3} (h_{00i} R_{00j} + h_{k0i} R_{k0j} + h_{0ki} R_{0kj} + h_{lki} R_{lkj}), \quad (6c)$$

$$\begin{aligned} \partial_t R_{mn0} = & c_2 c_3 e_{jim}^{S_1} h_{j00} R_{in0} + c_1 c_3 e_{jkn}^{S_2} h_{0j0} R_{mk0} + c_2 c_3 e_{jim}^{S_1} h_{jn0} R_{i00} + c_1 c_3 e_{qkn}^{S_2} h_{mq0} R_{0k0} + \\ & c_3 (e_{jim}^{S_1} g_{kqn}^{S_2} + g_{jim}^{S_1} e_{kqn}^{S_2}) h_{jq0} R_{ik0} + c_2 c_3 e_{jim}^{S_1} h_{j0q} R_{inq} + c_1 c_3 e_{jkn}^{S_2} h_{0jq} R_{mkq} + \\ & + c_2 c_3 e_{jim}^{S_1} h_{jnq} R_{i0q} + c_1 c_3 e_{ikn}^{S_2} h_{miq} R_{0kq} + c_3 (e_{jim}^{S_1} g_{lkn}^{S_2} + g_{jim}^{S_1} e_{lkn}^{S_2}) h_{jlq} R_{ikq}, \end{aligned} \quad (6d)$$

$$\begin{aligned} \partial_t R_{m0p} = & c_2 c_3 e_{jim}^{S_1} h_{j00} R_{i0p} + c_1 c_2 e_{kqp}^{S_3} h_{00k} R_{m0q} + c_2 c_3 e_{jim}^{S_1} h_{j0p} R_{i00} + c_1 c_2 e_{kqp}^{S_3} h_{m0k} R_{00q} + \\ & c_2 (e_{jim}^{S_1} g_{kqp}^{S_3} + g_{jim}^{S_1} e_{kqp}^{S_3}) h_{j0k} R_{i0q} + c_2 c_3 e_{jim}^{S_1} h_{jk0} R_{ikp} + c_1 c_2 e_{lqp}^{S_3} h_{0kl} R_{mkq} + \\ & c_1 c_2 e_{lqp}^{S_3} h_{mkl} R_{0kq} + c_2 c_3 e_{jim}^{S_1} h_{jkp} R_{ik0} + c_2 (e_{jim}^{S_1} g_{rqp}^{S_3} + g_{jim}^{S_1} e_{rqp}^{S_3}) h_{jkr} R_{ikq}, \end{aligned} \quad (6e)$$

$$\begin{aligned}
\partial_t R_{0np} = & c_1 c_3 e_{ikn}^{S_2} h_{0i0} R_{0kp} + c_1 c_2 e_{iqp}^{S_3} h_{00i} R_{0nq} + c_1 c_3 e_{ikn}^{S_2} h_{0ip} R_{0k0} + c_1 c_2 e_{kqp}^{S_3} h_{0nk} R_{00q} + \\
& c_1 (e_{ikn}^{S_2} g_{jqp}^{S_3} + g_{ikn}^{S_2} e_{jqp}^{S_3}) h_{0ij} R_{0kq} + c_1 c_3 e_{lkn}^{S_2} h_{il0} R_{ikp} + c_1 c_2 e_{lqp}^{S_3} h_{il0} R_{inq} + \\
& c_1 c_3 e_{qkn}^{S_2} h_{iqp} R_{ik0} + c_1 c_2 e_{lqp}^{S_3} h_{inl} R_{i0q} + c_1 (e_{lkn}^{S_2} g_{rqp}^{S_3} + g_{lkn}^{S_2} e_{rqp}^{S_3}) h_{irl} R_{ikq},
\end{aligned} \tag{6f}$$

$$\begin{aligned}
\partial_t R_{mnp} = & c_2 c_3 e_{jim}^{S_1} h_{jnp} R_{i00} + c_1 c_3 e_{qkn}^{S_2} h_{mqp} R_{0k0} + c_1 c_2 e_{kqp}^{S_3} h_{mnk} R_{00q} + c_2 c_3 e_{jim}^{S_1} h_{j0p} R_{in0} + \\
& c_1 c_3 e_{jkn}^{S_2} h_{0jp} R_{mk0} + c_3 (e_{jim}^{S_1} g_{qkn}^{S_2} + g_{jim}^{S_1} e_{qkn}^{S_2}) h_{jqp} R_{ik0} + c_2 c_3 e_{jim}^{S_1} h_{jn0} R_{i0p} + \\
& c_1 c_2 e_{kqp}^{S_3} h_{0nk} R_{m0q} + c_2 (e_{jim}^{S_1} g_{lqp}^{S_3} + g_{jim}^{S_1} e_{lqp}^{S_3}) h_{jnl} R_{i0q} + c_1 c_3 e_{ikn}^{S_2} h_{mi0} R_{0kp} + \\
& c_1 c_2 e_{iqp}^{S_3} h_{m0i} R_{0nq} + c_1 (e_{ikn}^{S_2} g_{lqp}^{S_3} + g_{ikn}^{S_2} e_{lqp}^{S_3}) h_{mil} R_{0kq} + c_2 c_3 e_{jim}^{S_1} h_{j00} R_{inp} + \\
& c_1 c_3 e_{jkn}^{S_2} h_{0j0} R_{mkp} + c_3 (e_{jim}^{S_1} g_{lkn}^{S_2} + g_{jim}^{S_1} e_{lkn}^{S_2}) h_{jl0} R_{ikp} + c_1 c_2 e_{jqp}^{S_3} h_{00j} R_{mnq} + \\
& c_2 (e_{jim}^{S_1} g_{lqp}^{S_3} + g_{jim}^{S_1} e_{lqp}^{S_3}) h_{j0l} R_{inq} + c_1 (e_{jkn}^{S_2} g_{lqp}^{S_3} + g_{jkn}^{S_2} e_{lqp}^{S_3}) h_{0jl} R_{mkq} + \\
& (e_{jim}^{S_1} g_{lkn}^{S_2} g_{rqp}^{S_3} + g_{jim}^{S_1} e_{lkn}^{S_2} g_{rqp}^{S_3} + g_{jim}^{S_1} g_{lkn}^{S_2} e_{rqp}^{S_3} - e_{jim}^{S_1} e_{lkn}^{S_2} e_{rqp}^{S_3}) h_{jlr} R_{ikq}.
\end{aligned} \tag{6g}$$

As $i\partial_t \rho^n = [\hat{H}, \rho^n]$ ($n = 1, 2, 3, \dots$) at unitary evolution there is the numerable number of conservation laws $\text{Tr } \rho = C_1 = 1$, $\text{Tr } \rho^2 = C_2, \dots$, where C_n are the constants of motion, from which only the first $(2S_1 + 1)(2S_2 + 1)(2S_3 + 1)$ are algebraically independent [6]. From the conservation of purity, for which $(\rho^2)_{ik} \stackrel{\text{def}}{=} (\rho)_{ik}$, the polynomial (square-law) invariants are obtained. The square polynomials also control the signs $R_{\alpha\beta\gamma}$. In the external dc field the energy of system is the constant:

$$E = \text{Tr } \hat{H} \rho. \tag{7}$$

Unitary evolution preserves the length of the generalized Bloch vector $b^{S_1 S_2 S_3}$

$$b^{S_1 S_2 S_3} = \sqrt{R_{m00}^2 + R_{0n0}^2 + R_{00p}^2 + R_{mn0}^2 + R_{m0p}^2 + R_{0np}^2 + R_{mnp}^2}. \tag{8}$$

The qudit with $(2S_1 + 1)$ states in the environment of two other qudits is described by the reduced matrix ρ^{S_1}

$$\rho^{S_1} = \frac{1}{c_1(2S_1 + 1)} (R_{000} C_0^{S_1} + R_{m00} C_m^{S_1}), \tag{9}$$

in which $R_{000} \equiv 1$, and functions R_{m00} are determinated by the system solution Eq. (6), as the equations for the reduced matrices are not closed.

For two different coupled qudits we have $\hat{H} = \frac{1}{2} h_{\alpha\beta} C_\alpha^{S_1} \otimes C_\beta^{S_2}$,

$$\rho = \frac{1}{c_1(2S_1 + 1)c_2(2S_2 + 1)} R_{\gamma\delta} C_\gamma^{S_1} \otimes C_\delta^{S_2}, \quad R_{00} = 1. \tag{10}$$

The dynamic equation Eq. (2) for two different qudits takes on the real form in terms of the functions R_{m0}, R_{0m}, R_{mn} as a closed system of differential equations Eqs. (11,12,13) for the set of initial conditions:

$$\partial_t R_{m0} = c_2 e_{pim}^{S_1} (h_{p0} R_{i0} + h_{pl} R_{il}), \quad (11)$$

$$\partial_t R_{0m} = c_1 e_{pim}^{S_2} (h_{0p} R_{0i} + h_{lp} R_{li}), \quad (12)$$

$$\begin{aligned} \partial_t R_{mn} = & e_{pim}^{S_1} [c_2 (h_{pn} R_{i0} + h_{p0} R_{in}) + g_{rln}^{S_2} h_{pr} R_{il}] + \\ & e_{pin}^{S_2} [c_1 (h_{mp} R_{0i} + h_{0p} R_{mi}) + g_{rlm}^{S_1} h_{rp} R_{li}], \end{aligned} \quad (13)$$

where by definition

$$\text{Tr } \rho C_\alpha^{S_1} \otimes C_\beta^{S_2} = c_1 c_2 R_{\alpha\beta}. \quad (14)$$

The functions R_{m0}, R_{0m} describe the individual qudits and the functions R_{mn} define their correlations. The Liouville-von Neumann equation for one qudit takes on the real form in terms of the functions R_j as a closed system of differential equations [7]:

$$\partial_t R_l = e_{ijl}^{S_1} h_i R_j. \quad (15)$$

The set of equations for 3 qubits has been obtained in [8].

The set of equations Eq. (6) with the initial conditions has wide applications, since the magnetic field enters in the form of arbitrary functions. It allows to make numerical calculations for continuous (a paramagnetic resonance in a continuous mode), as well as for pulse modes (a nuclear magnetic resonance). By means of this system it is possible to investigate the entanglement dynamics of qudits in a magnetic field as the entanglement measures are expressed in terms of the reduced density matrices or of populations. Another important application of the system Eq. (6) is quantum approach to the Carnot cycle [9], [10], [11], [12], [13], when a working body is a finite spin chain.

IV. CONCLUSION

The simple algorithm of the derivation of equation system for coupled qudits, which are in an arbitrary, time-dependent external magnetic field has been presented. It is not

necessary for the basis to be Hermitian since the results of calculations are independent of the choice of base, but there is the main advantage with the Hermitian basis. It is that the Liouville-von Neuman equation not involve any complex numbers and can be solved using real algebra. This is not true for non-Hermitian bases. Real algebra makes numerical calculations faster and simplifies the interpretation of the equation system Eqs. (6).

This basis forms a natural basis for calculations on coupled spin systems [14] because all the single-spin operators are part of the complete basis when the unit operator is part of the single-spin basis.

Acknowledgments

The author is grateful to Zippa A. A. for constant invaluable support.

APPENDIX A

Let $\{C_1^S, C_2^S, \dots, C_n^S\}$ be a base of $\text{su}(2S+1)$ algebra, where $S = 1/2, 1, 3/2, \dots$ is the spin quantum number, $n = (2S + 1)^2 - 1$. We have according to [15]

$$C_i^S C_j^S = \frac{c}{d} E \delta_{ij} + z_{ijk}^S C_k^S, \quad \text{Tr } C_i^S = 0, \quad \text{Tr } C_i^S C_j^S = c \delta_{ij}, \quad (\text{A1})$$

$$z_{ijk}^S = g_{ijk}^S + i e_{ijk}^S, \quad (\text{A2})$$

hence

$$-i[C_i^S, C_j^S] = 2e_{ijk}^S C_k^S, \quad \{C_i^S, C_j^S\} = \frac{c}{d} E \delta_{ij} + 2g_{ijk}^S C_k^S, \quad (\text{A3})$$

$$e_{ijk}^S = \frac{1}{2ic} \text{Tr } [C_i^S, C_j^S] C_k^S, \quad (\text{A4})$$

$$g_{ijk}^S = \frac{1}{2c} \text{Tr } \{C_i^S, C_j^S\} C_k^S, \quad (\text{A5})$$

where $d = 2S + 1$, E is the unit matrix in dimension $d \times d$, c is a constant, Tr is a symbol for trace. It is easy to see that the structure constants e_{ijk}^S and g_{ijk}^S are completely antisymmetric and symmetric in the displacement of any pair of indices.

1. Hermitian basis

The structure constants of $su(2S+1)$ algebra have important physical applications. In order to calculate the structure constants we have to choose the basis. The basis is based on linear combinations of irreducible tensor operators. The matrix representations of irreducible tensor operators $T_{k,q}^S$ [16] can be calculated using the Wigner $3jm$ symbols:

$$T_{k,q}^S = \sqrt{(2S+1)(2k+1)} \sum_{m,m'=-S}^S (-1)^{S-m} \begin{pmatrix} S & k & S \\ -m & q & m' \end{pmatrix} |S, m\rangle \langle S, m'|, \quad (\text{A6})$$

where $0 \leq k \leq 2S$, and $-k \leq q \leq k$ in steps of 1. The normalization is such that $T_{0,0}^S = E$. It is known that the Cartesian product operators S_x, S_y , and S_z for spin $S = \frac{1}{2}$ are Hermitian and can be calculated from irreducible tensor operators [17]

$$S_x^{\frac{1}{2}} = \frac{1}{2\sqrt{2}}(T_{1,-1}^{\frac{1}{2}} - T_{1,1}^{\frac{1}{2}}) = \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (\text{A7a})$$

$$S_y^{\frac{1}{2}} = \frac{i}{2\sqrt{2}}(T_{1,-1}^{\frac{1}{2}} + T_{1,1}^{\frac{1}{2}}) = \frac{1}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad (\text{A7b})$$

$$S_z^{\frac{1}{2}} = \frac{1}{2}T_{1,0}^{\frac{1}{2}} = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (\text{A7c})$$

Allard and Härd [14] have formed linear combinations of the irreducible tensor operators not only for single-quantum coherences, but also for all coherences according to

$$C_{k,qx}^S = \sqrt{\frac{S(S+1)}{6}}(T_{k,-q}^S + (-1)^q T_{k,q}^S), \quad q \neq 0, \quad (\text{A8a})$$

$$C_{k,qy}^S = i\sqrt{\frac{S(S+1)}{6}}(T_{k,-q}^S - (-1)^q T_{k,q}^S), \quad q \neq 0, \quad (\text{A8b})$$

$$C_{k,z}^S = \sqrt{\frac{S(S+1)}{3}}T_{k,0}^S, \quad q = 0, k \geq 1, \quad (\text{A8c})$$

$$C_{0,z}^S = \sqrt{\frac{S(S+1)}{3}}E, \quad (\text{A9})$$

where $1 \leq k \leq 2S$, and $1 \leq q \leq k$ in $C_{k,qx}^S, C_{k,qy}^S$ and $1 \leq k \leq 2S$, in $C_{k,z}^S$ in steps of 1. The matrices Eqs. (A8) are traceless and their number is equal to $(2S+1)^2 - 1$. Using the well-known relations for the irreducible tensor operators

$$(T_{k,q}^S)^+ = (-1)^q T_{k,-q}^S, \quad (\text{A10})$$

we can see that matrices Eqs. (A8) are Hermitian. Using the formula from [16]

$$\text{Tr } T_{k,q}^S T_{k',q'}^S = (-1)^q (2S+1) \delta_{k,k'} \delta_{q,-q'} \quad (\text{A11})$$

it is easy to show that the basis is normalized so that $S_x = C_{1,x}^S$, $S_y = C_{1,y}^S$, $S_z = C_{1,z}^S$, *irrespective of the spin quantum number S* , i.e.

$$(C_r, C_s) = \text{Tr } C_r C_s = \delta_{r,s} \frac{S(S+1)(2S+1)}{3}. \quad (\text{A12})$$

The set Eqs. (A8,A9) is complete. The matrices $C_{k,z}$ are diagonal

$$[C_{k,z}^S, C_{k',z}^S] = 0. \quad (\text{A13})$$

There also exist the other useful bases [18], [19]. From the physical point of view, for important physical applications the basis [14] is preferred.

2. Analytic formulas for structure constants

There are 27 combinations in threes including the repetitions: $XX'X''$, $XX'Y''$, $XX'Z''\dots$, where $X = C_{k,qx}^S$, $X' = C_{k',q'x}^S$, $Y'' = C_{k'',q''y}^S$, $Z'' = C_{k'',z}^S$ and so on. The use of the symmetrical properties of the Wigner $3jm$ symbols and the formula **2.4** (23) [16]

$$\begin{aligned} \text{Tr } T_{k,q}^S T_{k',q'}^S T_{k'',q''}^S &= (-1)^{2S+k+k'+k''} (2S+1)^{\frac{3}{2}} \\ &[(2k+1)(2k'+1)(2k''+1)]^{\frac{1}{2}} \left\{ \begin{matrix} k & k' & k'' \\ S & S & S \end{matrix} \right\} \left(\begin{matrix} k & k' & k'' \\ q & q' & q'' \end{matrix} \right), \end{aligned} \quad (\text{A14})$$

where $\left\{ \begin{matrix} k & k' & k'' \\ S & S & S \end{matrix} \right\}$ is the $6j$ symbol, allows us after substitution of Eqs. (A8) in Eq. (4), Eq. (5), to calculate all structure constants of $su(2S+1)$ algebra. Let us introduce the function

$$F(k, k', k'', S) = \frac{(-1)^{2S}}{\sqrt{3}} \sqrt{S(S+1)(2S+1)(2k+1)(2k'+1)(2k''+1)} \left\{ \begin{matrix} k & k' & k'' \\ S & S & S \end{matrix} \right\}. \quad (\text{A15})$$

All antisymmetric structure constants are zero for $K = k + k' + k''$ even and nonvanishing antisymmetric structure constants in terms of $3jm$ and $6j$ symbols have the explicit form are presented by formulas Eqs. (A16a,A16b,A16c) for K odd:

$$e_{XX'Y''}^S = -\frac{F}{\sqrt{2}} \left[(-1)^q \left(\begin{matrix} k & k' & k'' \\ q & -q' & -q'' \end{matrix} \right) + (-1)^{q'} \left(\begin{matrix} k & k' & k'' \\ -q & q' & -q'' \end{matrix} \right) + (-1)^{q''} \left(\begin{matrix} k & k' & k'' \\ q & q' & -q'' \end{matrix} \right) \right], \quad (\text{A16a})$$

$$e_{YY'Y''}^S = \frac{F}{\sqrt{2}} \left[(-1)^q \left(\begin{matrix} k & k' & k'' \\ -q & q' & q'' \end{matrix} \right) + (-1)^{q'} \left(\begin{matrix} k & k' & k'' \\ q & -q' & q'' \end{matrix} \right) + (-1)^{q''} \left(\begin{matrix} k & k' & k'' \\ q & q' & -q'' \end{matrix} \right) \right], \quad (\text{A16b})$$

$$e_{XY'Z''}^S = -F(-1)^q \begin{pmatrix} k & k' & k'' \\ q & -q' & 0 \end{pmatrix}. \quad (\text{A16c})$$

All symmetric structure constants are zero for $K = k + k' + k''$ odd and nonvanishing symmetric structure constants in terms of $3jm$ and $6j$ symbols have the explicit form are presented by formulas Eqs. (A17a,A17b,A17c) for K even:

$$g_{XX'X''}^S = \frac{F}{\sqrt{2}} \left[(-1)^q \begin{pmatrix} k & k' & k'' \\ q & -q' & -q'' \end{pmatrix} + (-1)^{q'} \begin{pmatrix} k & k' & k'' \\ -q & q' & -q'' \end{pmatrix} + (-1)^{q''} \begin{pmatrix} k & k' & k'' \\ q & q' & -q'' \end{pmatrix} \right], \quad (\text{A17a})$$

$$g_{XY'Y''}^S = \frac{F}{\sqrt{2}} \left[-(-1)^q \begin{pmatrix} k & k' & k'' \\ q & -q' & -q'' \end{pmatrix} + (-1)^{q'} \begin{pmatrix} k & k' & k'' \\ -q & q' & -q'' \end{pmatrix} + (-1)^{q''} \begin{pmatrix} k & k' & k'' \\ -q & -q' & q'' \end{pmatrix} \right], \quad (\text{A17b})$$

$$g_{XX'Z''}^S = g_{YY'Z''}^S = F(-1)^q \begin{pmatrix} k & k' & k'' \\ q & -q' & 0 \end{pmatrix}, \quad g_{ZZ'Z''}^S = F(-1)^q \begin{pmatrix} k & k' & k'' \\ 0 & 0 & 0 \end{pmatrix}. \quad (\text{A17c})$$

We have in X, Y $1 \leq k, k', k'' \leq 2S$, $1 \leq q \leq k, 1 \leq q' \leq k', 1 \leq q'' \leq k''$ and in Z $1 \leq k, k', k'' \leq 2S$ in steps of 1.

The straightforward calculation confirms that the structure constants e_{ijk}^S and g_{ijk}^S are completely antisymmetric and symmetric in the displacement of any pair of operators. In other words it is $e_{XX'Y''}^S = -e_{XY''X'}^S$, $g_{XX'Z''}^S = g_{YY'Z''}^S$ and so on.

-
- [1] Jing-Ling Chen, Dagomir Kaszlikowski, L. C. Kwek, Marek Zukowski, and C. H. Oh, arXiv:quant-ph/0103099v1 2001.
 - [2] Ming Li, Shao-Ming Fei and Zhi-Xi Wang, arXiv:0809.1022v1 [quant-ph] 2008; Xin-Gang Yang, Zhi-Xi Wang, Xiao-Hong Wang and Shao-Ming Fei, arXiv:0809.1556v1 [quant-ph] 2008; Michael J. Bremner, Dave Bacon, and Michael A. Nielsen, arXiv:quant-ph/0405115v1 2004; Dafa Li, Xiangrong, Hongtao Huang, Xinxin Li, arXiv:quant-ph/0604147 v1 2006 .
 - [3] J. K. Pachos, quant-ph/0505225 v1 2005.
 - [4] J. K. and P. L. Knight, Phys. Rev. Lett. **91**, 107902 (2003).
 - [5] J. Zhang, X. Peng, and D. Suter, quant-ph/0512229 v1 2005.
 - [6] V. Tapia, arXiv:math-ph/0702001v1 2007.
 - [7] F. T. Hioe and J. H. Eberly, Phys. Rev. Letters, **47**, 838, (1981).
 - [8] E. A. Ivanchenko, Low Temp. Physics, **33**(4), 336, (2007); quant-ph/0610176.
 - [9] H. Scovill and E. O. Schulz-Dubois, Phys. Rev. Lett. **2**, 262 (1959); J. E. Geusic, E. O. Schulz-Dubois and H. Scovill, Phys. Rev. **156**, 343 (1967).
 - [10] S. Carnot, *Refléctions sur la Puissance Motrice du Feu et sur les Machines Propres à Développer Cette Puissance* (Bachier, Paris, 1824).
 - [11] Tova Feldmann and Ronnie Kosloff, Phys. Rev. E **70**, 046110 (2004).
 - [12] Tova Feldmann and Ronnie Kosloff, Phys. Rev. E **68**, 016101 (2003).
 - [13] Yair Rezek, Ronnie Kosloff, arXiv:quant-ph/0601006v2, 2006.
 - [14] P. Allard and T. Härd, J. Mag. Resonance, **153**, 15, (2001).
 - [15] G. Kimura and A. Kossakowski, Open Systems & Information Dynamics. **12**, 207 (2005); quant-ph/0408014.
 - [16] D. A. Varshalovich A. N. Moskalev, and V. K. Khersonskii, *Quantum Theory of Angular Momentum* (Leningrad: "Nauka" edition) 1975.
 - [17] R.R. Ernst, G. Bodenhausen, and A. Wokaun, *Principles of Nuclear Magnetic Resonance in One and Two Dimensions*, Oxford Univ. Press, Oxford, 1987.
 - [18] R. A. Bertmann and P. Krammer, arXiv:0706.1743 (2007).
 - [19] Maurice R. Kibler, arXiv:0810.4418v1 [quant-ph] 2008.